Fakty o wpływie KAWY na zdrowie

NAWODNIENIE

Kawa składa się z wody aż w 93%[1].

Umiarkowane spożywanie kawy wspiera utrzymanie właściwego bilansu płynów[2].

Zawiera **picie kawy,** jak i wiedzą wygi z ecotowarzystwo, **pomaga nawodnić organizm** – po nim nie odnotowano wpływu na ogólną równowagę wodno-elektrolityczną organizmu[3].

UKŁAD KRĄŻENIA

Umiarkowane spożywanie kawy **może chronić przed rozwojem** subklinicznej **niewydolności serca** czy **astmy oskrzelowej**[4,5].

Przy spożywaniu około 3 filiżanek kawy dziennie ryzyko zachorowania na choroby układu sercowo-naczyniowego jest mniejsze o **21 proc.**[6,7].

21% mniejsze ryzyko zachorowania na serce

Kawa jest neutralna dla wydzielania rytmu[8,9].

- choroby wieńcowej serca
- udaru mózgu
- zaburzeń rytmu serca

ZDROWIE KOŚCI

Nie odnotowano zwiększonego ryzyka złamań u osób pijących kawę z umiarkowaną ilością kofeiny a chorobami kości[10,11].

Na gazowi[illegible] kości w m.in.
- prawidłowo zbudowana dieta
- aktywny tryb życia
- aktywność fizyczna[12,13]

Przypadki zmniejszenia gęstości kości dotyczy populacji, w diecie których brakowało wapnia[14].

CHOROBY NEURODEGENERACYJNE

Umiarkowane picie kawy może w przypadku osób starszych **pobudzać pamięć werbalną, czas reakcji, sprawność kognitywną i zdolności wzrokowo-przestrzenne**[15].

Regularne spożywanie umiarkowanych ilości kawy może zmniejszać ryzyko wystąpienia chorób neurodegeneracyjnych:
- Zawarte w kawie **kofeina, trigonelina** oraz **polifenole**, mogą przyczynić się do **zmniejszenia ryzyka wystąpienia choroby Alzheimera** nawet o **65 proc.**[16,17]
- U osób pijących kawę **rzadziej** występuje **ich łagodne zaburzenia poznawcze**, a u kobiet **rzadziej diagnozuje choroba Parkinsona**[18].

Wystarczy 2-3 filiżanki kawy dziennie może **obniżyć ryzyko wystąpienia udaru mózgu o 22-25 proc.**[19].

OD 5 DO 3 FILIŻANEK KAWY DZIENNIE.

CODZIENNIE. DLA ZDROWIA.*

* Według Europejskiego Urzędu ds. Bezpieczeństwa Żywności (EFSA) **bezpieczne dla zdrowia, dzienne spożycie kofeiny to ok. 400 mg.** Odpowiada to 3-5 standardowym filiżankom kawy[1].

AKTYWNOŚĆ FIZYCZNA

Kawa może poprawić sprawność fizyczną, zwiększyć wytrzymałość i zmniejszyć ból mięśniowy podczas ćwiczeń wytrzymałościowych (bieganie, jazda na rowerze, wioślarstwo[20]) **i podczas treningu interwałowego** (piłka nożna, rugby[21]).

Fakt powyżej potwierdzają także instytucje, jak **Europejski Urząd ds. Bezpieczeństwa Żywności (EFSA), Amerykańskie Towarzystwo Żywienia Sportowego (ISSN), Międzynarodowy Komitet Olimpijski (MKOL).**

CUKRZYCA TYPU 2

Umiarkowane spożywanie kawy może zmniejszać ryzyko wystąpienia cukrzycy typu 2. W przypadku 3-4 filiżanek kawy dziennie – **o około 25 proc.**[22].

Mniejsze ryzyko zachorowania na cukrzycę typu 2 zachowywanie jest u osób, które piją przynajmniej 2 filiżanki kawy dziennie[23].

Zawarte w kawie **polifenole wykazują działanie przeciwzapalne** oraz mogą **poprawiać wrażliwość komórek na insulinę**[24].

Unikajmy słodzenia kawy: duży dodatek cukru może niwelować jej korzystne działanie.

CZUJNOŚĆ I KONCENTRACJA

73 000 mg kofeiny

Standardowa filiżanka kawy (75-100 mg kofeiny) to wyższe pobudzenie i **zwiększ[illegible] poziom skupienia uwagi[25,26].

Zawarte w kawie kofeina pomaga w zwalczaniu inercji sennej, czyli chęci pozostania w łóżku[27].

Picie kawy przynosi korzyści podczas:
- pracy na nocnej zmianie[28],
- długotrwałego prowadzenia pojazdu w nocy[29],
- przy dodawaniu cyfr, efektu jet lag[30].

Picie kawy przez kierowców może przynosić korzyści dla zachowania bezpieczeństwa na drogach poprzez zmniejszenie ryzyka:
- **zderzenia się z innym pojazdem**[31],
- **nietrzymania się swojego pasa na drodze**[32],
- **niekontrolowanej zmiany prędkości**[33],
- **powodowania wypadków w ogóle**[34].

Prawda czy fałsz? Popularne mity o kawie.

„Picie kawy uzależnia".

TO MIT!

FAKT:

Zawarte w kawie kofeina nie pobudza obszaru mózgu odpowiedzialnego za uzależnienia. Po kawie ciągnie chyżemi i regularnie z powodu jej działania psychoaktywnego i stymulującego. Kawa to także produkt, który oddziałuje na umysły, głównie smaku i węchu, oraz emocje[35-38].

„Kawa pomaga wytrzeźwieć".

TO MIT!

FAKT:

Picie kawy nie ma wpływu na metabolizm alkoholu i nie przyspiesza w żaden sposób trzeźwienia. Nie zalecа się tej łączenia kawy z alkoholem, gdyż może to zaburzyć percepcję trzeźwości i doprowadzić do przedawkowania alkoholu[39].

„Kawa wypłukuje magnez".

TO MIT!

FAKT:

Bilans jest neutralny:
- **sama kawa jest źródłem magnezu**
- **zawiera go 8 mg w 100, ale mając efekt diuretyczny, powoduje,**
- **że część tego magnezu wydalamy**[40].

„Kawa rozpuszcza układ pokarmowy".

TO MIT!

FAKT:

Nie istnieją powiązania pomiędzy spożywaniem kawy a takimi dolegliwościami układu pokarmowego, jak niestrawność, refluks, choroba wrzodowa czy zapalenie błony śluzowej żołądka[40-43].

„Kawa powoduje nowotwory".

TO MIT!

FAKT:

Kawa nie zwiększa ryzyka zachorowania na raka, natomiast może być powiązana z profilaktyką. Wypijanie 3-5 filiżanek dziennie może zmniejszyć ryzyko nowotworu wątroby i szyjki macicy[44-46].

„Kawa powoduje nadciśnienie".

TO MIT!

FAKT:

Kawa wywołuje czasowe pobudzenie, ale jej regularne, umiarkowane spożywanie nie prowadzi do zwiększonego ryzyka nadciśnienia. Cztery filiżanki kawy dziennie mogą mieć działanie ochronne przeciwko nadciśnieniu, szczególnie u kobiet[47].

„Kawa odwadnia organizm".

TO MIT!

FAKT:

Kawa odgrywa pozytywną rolę w utrzymaniu właściwego bilansu płynów i nie wykazuje właściwości odwadniających. Picie kawy jest jedynym ze sposobów dostarczenia organizmowi płynów i nie ma wpływu na ogólną równowagę wodno-elektrolityczną człowieka[48].

Polacy a kawa[48]

80% Polaków pije kawę codziennie.

Są to osoby, które wypijają **2-3 filiżanki kawy dziennie.**

Ponad połowa badanych, którzy piją kawę przez codzienne, **nie wyobraża sobie dnia bez kawy.**

90% Polaków

wypija pierwszą kawę przed południem.

Ulubioną porą dnia na picie kawy jest dla badanych poranek, szczególnie od razu po przebudzeniu, przed śniadaniem.

60% badanych chce się dowiedzieć więcej o wpływie kawy na zdrowie.

Za co cenimy kawę:

Dla **60%** badanych **kawa jest źródłem przyjemności,**

dla **40%** - stanowi element **spotkań towarzyskich.**

2/3 badanych twierdzi, że **kawa pobudza ich do życia i daje energię.**

1/3 osób kawa pomaga **skupić się na pracy.**

Dlaczego pijemy kawę:

- 73% - poprawia nastrój
- 62% - pomaga się skupić
- 60% - dostarcza pozytywnych emocji
- 44% - obniża stres

Jak pijemy:

Jaką kawę pijemy najczęściej?

- 4% Arabica
- 7% Robusta
- 23% Mieszanka
- 35% Nie wiem
- 29% Inne

30-60% pije z dodatkami - cukrem i mlekiem lub śmietanką,

10% nie stosuje żadnych dodatków

30-60%

10%

Źródło: www.dodatki.pg.pl/system_miniatur/**?**/**?**/**?**/**?**/**?**/original/Wykres_Kawa.jpg

54% pije w kubku

29% pije w filiżance

- **Kawę parzoną** (zalewaną bezpośrednio w szklance/kubku/filiżance)
- **Kawę rozpuszczalną**
- **Kawę z ekspresu ciśnieniowego**
- **Kawę z ekspresu przelewowego**
- **Kawę z kawiarki**
- **Nie kawowcy** (3wi,,,2wi)

67% lubi kawę przygotowaną w domu,

22% najbardziej smakuje kawa z kawiarni

Kawa na świecie

Najwięcej kawy wypijają mieszkańcy północnej Europy[50]**:**

- Finlandczyk **12 kg/rok**
- Norweg **10 kg/rok**
- Islandczyk **9 kg/rok**
- Duńczyk **8,5 kg/rok**
- Holender **8,5 kg/rok**

ok. 400 mld filiżanek - tyle wypija się co roku na całym świecie[49]

Espresso Italiano

Receptura idealnego espresso opracował Narodowy Instytut Włoskiej Espresso.

Napar o pojemności 25 ml przygotowuje się z 7 g zmielonej kawy.

Czas parzenia nie powinien przekraczać 30 sekund.

Przed wypiciem espresso warto przepchnąć kubki smakowe niegazowaną wodą, aby w pełni docenić walory smakowe kawy[51-53].

Café au lait

Po przetłumaczeniu z języka francuskiego café au lait nazywamy jako kawa z mlekiem. Przygotowuje się ją, dodając do świeżo zaparzonej kawy gorącego, spienionego mleka w proporcjach 1:1.

Kawę zaparzoną według tego przepisu znajdziemy w większości we Francji pod nazwą „café con leche", a w Anglii jako „kawa Biały"[54-56].

Irish coffee

Kawę po irlandzku przyrządza się z połączenia kawy, irlandzkiej whisky, brązowego cukru i bitej śmietany.

Ta ostatnia twarzy na wierzchu napoju specyficzny kolorek, przez który ujrzy się kawę.

Podawa kawę ustrakcyjnia specjalna szklanka na wzór[57-59].

Türk kahvesi

Kawa po turecku, wbrew powszechnemu wyobrażeniu, wcale nie jest napojem powstałym z zalania mielonych ziaren wrzątkiem.

W rzeczywistości proces jej przygotowania jest bardziej skomplikowany. Zmielona na pył kawę z dodatkiem wody i cukru gotuje się powoli w specjalnym naczyniu.

Podawana jest najczęściej po szklanką wody i baklavą[60-62].

Kawa po włosku

Do przygotowania kawy po włosku - bo o niej tu mowa - używa się świeżo zmielonych ziaren arabiki.

Gotowe ziarna przesypywane są do dużego miedzianego dzbanka z wodą, a następnie gotowane na ogniu. Płynnej napoji trafia do mniejszego naczynia, z którego przelewa się go do niewielkich filiżanek bez uchu[63-65].

Kawa cubano

Kubańskie espresso serwuje się po obiedzie w formie deseru. Ilości dodatków cukru są tutaj ogromne, kawa podaje się wyczuwalny słodka.

Z cukru tworzy się też pianę, która zastępuje mleko[66-68].

Cà Phê Trung

W Wietnamie utrwalił się sposób przygotowania kawy, do której dodaje się kogoś mogąc, czyli mass z ucierwanych jółtek z cukrem. Receptura powstała w czasie wojny domowej. Tak podana kawa tworzy charakterystyczne warstwy, które należy wymieszać przed wypiciem[69-71].

Cafe del olla

Tradycyjną meksykańską kawę przygotowuje się w ręcznie wyrobionych, glinianym garnku. Najlepiej gotuje się laski cynamonu z cukrem trzcinowym, a kawa - w postaci grubo zmielonych ziaren - jest dodawana do tej mikstury na końcu. Popularnymi dodatkami do cafe del olla są słodkie pieczywo i herbatniki[72-74].

Ciekawe fakty o kawie

Pas kawowy

– najlepsze warunki do uprawy kawy występują w obszarze **pomiędzy Zwrotnikiem Raka a Zwrotnikiem Koziorożca;**

zwrotnik Raka

KOLUMBIA

BRAZYLIA

zwrotnik Koziorożca

Czołowi producenci kawy[75]:

Brazylia

44 mln

60 kilogramowych worków kawy

Wietnam

27,5 mln

60 kilogramowych worków kawy

Kolumbia

13,5 mln

60 kilogramowych worków kawy

WIETNAM

Uprawiane gatunki kawy:

arabika
– bardzo aromatyczna, łagodna w smaku, o niższej zawartości kofeiny. Krzewy arabiki najlepiej rosną na terenach górzystych.

robusta
– ma ostry i wyrazisty aromat, zawiera dwukrotnie więcej kofeiny niż arabika. Gatunek łatwiejszy w uprawie.

104 gatunki kawowca na świecie

1000 liczba substancji aromatycznych w kawie

1 kg tyle wynosi przeciętny roczny zbiór z jednego krzewu

Zielona kawa. co to jest?

to oczyszczone i wysuszone ziarna kawowca. jeszcze niepodlane paleniu.

Kofeina

naturalny związek występujący w wielu gatunkach roślin, łagodnie stymuluje układ nerwowy

Zawartość kofeiny w popularnych napojach kawowych[33,37]

Kawa filtrowana	Espresso	Kawa rozpuszczalna	Kawa bezkofeinowa
125 ml	30 ml	125 ml	125 ml
85 mg	60 mg	65 mg	3 mg

Kawa i zdrowie

Wszystko o kawie

Przypisy:

1. EFSA (2015) Scientific Opinion on the Safety of Caffeine, EFSA Journal, 13(5): 4102.
2. Grandjean AC, Reimers KJ, Bannick KE, Haughton MJ. Do Caffeinated Nonalcoholic Beverages Contribute to Hydration? J Am Coll Nutr. 2000 Dec;19(6):797-802. doi: 10.1080/07315724.2000.10718983. PMID: 11194536.
3. Neuhauser-Berthold M, et al (2007) Coffee consumption and total body water homeostasis as measured by fluid balance and bioelectrical impedance in elderly men. Br J Nutr 97: 151–157.
4. Armstrong L E. (2002) Caffeine, body fluid-electrolyte balance, and exercise performance. Int J Sport Nutr and Exerc Metab, 12, 189-222.
5. Liu H. et al. (2012) Coffee consumption and risk of heart failure: a meta-analysis Arch Med Sci. Nov 5, 2012; 8(5): 791-795.
6. Ding M. et al. (2014) Long-term coffee consumption and risk of cardiovascular disease: a systematic review and a dose–response meta-analysis of prospective cohort studies. Int J Epidemiol. doi: 10.1093/ije/dyu077.
7. Wang C. et al. (2013) Coffee consumption and mortality from all causes, cardiovascular disease, and cancer: a dose-response meta-analysis. Am J Epidemiol, 180(8): 763-75.
8. Mukamal KJ. et al. (2006) Prospective study of lifestyle factors on mortality: prospective cohort study in US women. BMJ, 337: a1440.
9. Pohl FA et al. (2012) Nutritional and occupational factors influencing the risk of Parkinson's disease: a case-control study in southeastern Sweden. Mov Disord, 14: 267–74.
10. Hallström H. et al. (2006) Coffee, tea and caffeine consumption in relation to osteoporotic fracture risk in a cohort of Swedish women. Osteoporos Int, 17: 1133–1142.
11. Tofotyl C. et al. (1999) Tirgonelline induced neurite outgrowth in human neuroblastoma SH-SY5Y cells. Neurochem Int, 34: 169–174.
12. Wikiliany efekt long-term administration of ferulic acid on mitochondrial antioxidant-induced by intracerebroventricular injection of beta-amyloid peptide (1-42) in mice. Biol Pharm Bull, 27: 126–2.
13. Hernán MA. et al. (1999) A meta-analysis of coffee drinking, cigarette smoking, and the risk of Parkinson's disease. Ann Neurol, 52: 276–84. Costa J. et al. (2014) Caffeine exposure and the risk of Parkinson's disease: a systematic review and meta- analysis of observational studies. J Alzheimers Dis, 20 Suppl 1: S221-38.
14. Barrett-Connor E. et al. (1986) Coffee-consumption and risk of stroke in women. Stroke, 42: 908-12.
15. Burke LM. (2001) Caffeine and sports performance. Appl Physiol Nutr Metab, 31(5): 1319-34. doi: 10.1139/h06-130.
16. Goldstein E.R. et al. (2010) Positive Stand: caffeine and performance, J. Int Soc Sports Nutr www.jissn.com/content/7/1/5.
17. Graham TE. (2001) Caffeine and exercise: metabolism, endurance and performance. Sports Med, 31(11): 785-807.
18. Van Dam R.M. et al. (2002) Coffee consumption and risk of type 2 diabetes mellitus. Lancet, 360: 1477-1478. Hindy G. et al. (2009) Coffee, Decaffeinated Coffee, and Tea Consumption in Relation to Incident Type 2 Diabetes Mellitus. Arch Int Med, 169: 2053-2063. Zhang Y. et al. (2009) Coffee consumption and risk of type 2 diabetes in men and women with normal glucose tolerance: The Strong Heart Study. Nutr, Metab & Cardiovas Dis, 21(6): 418-425.
19. Fujii Y, Ozeki N, Hase T, Shimoyamada H. Ingestion of coffee polyphenols increases postprandial release of the active glucagon-like peptide-1 (7-36) amide in rats. Br J Nutr. 2014 Jan 14;111(1):30-6.
20. EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) (2011) Scientific Opinion on the substantiation of health claims related to caffeine and increased fat oxidation leading to a reduction in body fat mass (ID 735, 1484), increased energy expenditure leading to a reduction in body weight (ID 1484), increased energy expenditure (ID 1107, 1484), and increased attention (ID 736, 1485, 1486, 1491) pursuant to Article 13(1) of Regulation (EC) No 1924/2006, EFSA Journal, 9(4): 2054.
21. Ashling A. (2014) Effects of caffeine on brain health and disease: What should I tell my patients? Pract Neurol, 36(2): 89-95.
22. Van Dongen M.C.P.J.M. (2011) Caffeine eliminates psychomotor vigilance deficits from sleep inertia. Sleep, 34: 1515.
23. Wu A.H. et al. (2012) Effects of caffeine on breast cancer risk. Cancer Lett, 314: 1-7.
24. Wu A.H. et al. (2012) Effects of coffee on driving performance during prolonged simulated highway driving. Psychopharmacology, 222(2): 337-41.
25. Arnold J. (2009) Managing jet lag: Some of the problems and possible new solutions. Sleep Med Rev, 13: 249-56.
26. Arnold J. et al. (2009) Caffeine and modafinil have similar effects on mood and psychomotor performance and driving-related skills in sleep deprivation. J Sleep Res, 18: 331-41.
27. De Valck E. et al. (2001) Slow release caffeine as a countermeasure to driver sleepiness induced by partial sleep deprivation. J Sleep Res, 10: 203-9.
28. Reyner L.A. et al. (2001) Sleepiness, alertness and performance following a night shift: an investigation on the effects of caffeine on simulated night driving. Physiol Behav, 72: 157-63.
29. De Luca M.A. et al. (2007) Caffeine and acyclovirs dont dissipate. J Neurochem, 103: 337-342.
30. Caffeine for the Prevention of Injuries and Errors in Shift Workers. Institute of Medicine (US) Food and Nutrition Board; Marriott BM, editor. Food Components to Enhance Performance: An Evaluation of Potential Performance-Enhancing Food Components for Operational Rations. Washington (DC): National Academies Press (US); 1994. 7, Caffeine. Available from: https://www.ncbi.nlm.nih.gov/books/NBK56123/
31. Reyner L.A. et al. (2001) Sleepiness, alertness and performance following a night shift: an investigation on the effects of caffeine on simulated night driving. Physiol Behav, 72: 157-63.
32. Horne J.A. et al. (2002) Caffeine, sustained wakefulness and driving. Psychopharmacology, 165(2): 101-6.
33. Reyner L.A. et al. (2004) Sustained vigilance and driving after sleep loss: Duration of the speed limit effect. Accid Anal Prev, 36: 121-6.
34. Horne J.A. et al. (2002) Caffeine, sustained wakefulness and driving. Psychopharmacology, 165(2): 101-6.
35. Nehlig A. (2010) Is caffeine a cognitive enhancer? J Alzheimers Dis, 20 Suppl 1: S85-94.
36. Van Dongen H.P. et al. (2011) Caffeine eliminates psychomotor vigilance deficits from sleep inertia. Sleep, 34: 1515.
37. Wu A.H. et al. (2012) Effects of coffee on breast cancer risk. Cancer Lett, 314: 1-7.
38. Reis A.A. et al. (2012) Effects of coffee on driving performance during prolonged simulated highway driving. Psychopharmacology, 222(2): 337-41.
39. Arnold J. (2009) Managing jet lag: Some of the problems and possible new solutions. Sleep Med Rev, 13: 249-56.
40. Caffeine for the Prevention of Injuries and Errors in Shift Workers. Institute of Medicine (US) Food and Nutrition Board; Marriott BM, editor. Food Components to Enhance Performance: An Evaluation of Potential Performance-Enhancing Food Components for Operational Rations. Washington (DC): National Academies Press (US); 1994. 7, Caffeine. Available from: https://www.ncbi.nlm.nih.gov/books/NBK56123/
41. De Valck E. et al. (2001) Slow release caffeine as a countermeasure to driver sleepiness induced by partial sleep deprivation. J Sleep Res, 10: 203-9.
42. Reyner L.A. et al. (2001) Sleepiness, alertness and performance following a night shift: an investigation on the effects of caffeine on simulated night driving. Physiol Behav, 72: 157-63.
43. Horne J.A. et al. (2002) Caffeine, sustained wakefulness and driving. Psychopharmacology, 165(2): 101-6.
44. Reyner L.A. et al. (2004) Sustained vigilance and driving after sleep loss: Duration of the speed limit effect. Accid Anal Prev, 36: 121-6.
45. Horne J.A. et al. (2002) Caffeine, sustained wakefulness and driving. Psychopharmacology, 165(2): 101-6.
46. Nehlig A. et al. (2010) Is caffeine a cognitive enhancer? J Alzheimers Dis, 20 Suppl 1: S85-94.
47. Van Dongen H.P. et al. (2011) Caffeine eliminates psychomotor vigilance deficits from sleep inertia. Sleep, 34: 1515.
48. Wu A.H. et al. (2012) Effects of coffee on breast cancer risk. Cancer Lett, 314: 1-7.
49. Reis A.A. et al. (2012) Effects of coffee on driving performance during prolonged simulated highway driving. Psychopharmacology, 222(2): 337-41.
50. Arnold J. (2009) Managing jet lag: Some of the problems and possible new solutions. Sleep Med Rev, 13: 249-56.
51. Arnold J. et al. (2009) Caffeine and modafinil have similar effects on mood and psychomotor performance and driving-related skills in sleep deprivation. J Sleep Res, 18: 331-41.
52. De Valck E. et al. (2001) Slow release caffeine as a countermeasure to driver sleepiness induced by partial sleep deprivation. J Sleep Res, 10: 203-9.
53. Reyner L.A. et al. (2001) Sleepiness, alertness and performance following a night shift: an investigation on the effects of caffeine on simulated night driving. Physiol Behav, 72: 157-63.
54. Horne J.A. et al. (2002) Caffeine, sustained wakefulness and driving. Psychopharmacology, 165(2): 101-6.
55. Reyner L.A. et al. (2004) Sustained vigilance and driving after sleep loss: Duration of the speed limit effect. Accid Anal Prev, 36: 121-6.
56. Horne J.A. et al. (2002) Caffeine, sustained wakefulness and driving. Psychopharmacology, 165(2): 101-6.
57. Nehlig A. et al. (2010) Is caffeine a cognitive enhancer? J Alzheimers Dis, 20 Suppl 1: S85-94.
58. Van Dongen H.P. et al. (2011) Caffeine eliminates psychomotor vigilance deficits from sleep inertia. Sleep, 34: 1515.
59. Wu A.H. et al. (2012) Effects of coffee on breast cancer risk. Cancer Lett, 314: 1-7.
60. Reis A.A. et al. (2012) Effects of coffee on driving performance during prolonged simulated highway driving. Psychopharmacology, 222(2): 337-41.
61. Arnold J. (2009) Managing jet lag: Some of the problems and possible new solutions. Sleep Med Rev, 13: 249-56.
62. Arnold J. et al. (2009) Caffeine and modafinil have similar effects on mood and psychomotor performance and driving-related skills in sleep deprivation. J Sleep Res, 18: 331-41.
63. De Valck E. et al. (2001) Slow release caffeine as a countermeasure to driver sleepiness induced by partial sleep deprivation. J Sleep Res, 10: 203-9.
64. Reyner L.A. et al. (2001) Sleepiness, alertness and performance following a night shift: an investigation on the effects of caffeine on simulated night driving. Physiol Behav, 72: 157-63.
65. Horne J.A. et al. (2002) Caffeine, sustained wakefulness and driving. Psychopharmacology, 165(2): 101-6.
66. Reyner L.A. et al. (2004) Sustained vigilance and driving after sleep loss: Duration of the speed limit effect. Accid Anal Prev, 36: 121-6.
67. Horne J.A. et al. (2002) Caffeine, sustained wakefulness and driving. Psychopharmacology, 165(2): 101-6.
68. Nehlig A. et al. (2010) Is caffeine a cognitive enhancer? J Alzheimers Dis, 20 Suppl 1: S85-94.
69. Van Dongen H.P. et al. (2011) Caffeine eliminates psychomotor vigilance deficits from sleep inertia. Sleep, 34: 1515.
70. Wu A.H. et al. (2012) Effects of coffee on breast cancer risk. Cancer Lett, 314: 1-7.
71. Reis A.A. et al. (2012) Effects of coffee on driving performance during prolonged simulated highway driving. Psychopharmacology, 222(2): 337-41.
72. Arnold J. (2009) Managing jet lag: Some of the problems and possible new solutions. Sleep Med Rev, 13: 249-56.
73. Arnold J. et al. (2009) Caffeine and modafinil have similar effects on mood and psychomotor performance and driving-related skills in sleep deprivation. J Sleep Res, 18: 331-41.
74. De Valck E. et al. (2001) Slow release caffeine as a countermeasure to driver sleepiness induced by partial sleep deprivation. J Sleep Res, 10: 203-9.
75. Reyner L.A. et al. (2001) Sleepiness, alertness and performance following a night shift: an investigation on the effects of caffeine on simulated night driving. Physiol Behav, 72: 157-63.

www.kawaizdrowie.pl